arXiv:1501.06417v1 [math.FA] 26 Jan 2015

WEAKLY ALMOST PERIODIC BANACH ALGEBRAS ON SEMI-GROUPS

BAHRAM KHODSIANI, ALI REJALI

Abstract. Let $\mathcal{WAP}(\mathcal{A})$ be the space of all weakly almost periodic functionals on a Banach algebra $\mathcal{A}$. The Banach algebra $\mathcal{A}$ for which the natural embedding of $\mathcal{A}$ into $\mathcal{WAP}(\mathcal{A})^*$ is bounded below is called a $\mathcal{WAP}$-algebra. We show that the second dual of a Banach algebra $\mathcal{A}$ is a $\mathcal{WAP}$-algebra, under each Arens products, if and only if $\mathcal{A}^{**}$ is a dual Banach algebra. This is equivalent to the Arens regularity of $\mathcal{A}$. For a locally compact foundation semigroup S, we show that the absolutely continuous semigroup measure algebra $M_a(S)$ is a $\mathcal{WAP}$-algebra if and only if the measure algebra $M_b(S)$ is so.

1. Introduction and Preliminaries

The dual $\mathcal{A}^*$ of a Banach algebra $\mathcal{A}$ can be turned into a Banach $\mathcal{A}-$module in a natural way, by setting

$$\langle f\cdot a, b\rangle = \langle f, ab\rangle \qquad \text{and} \qquad \langle a\cdot f, b\rangle = \langle f, ba\rangle \qquad (a, b\in\mathcal{A}, f\in\mathcal{A}^*).$$

A *dual Banach algebra* is a Banach algebra $\mathcal{A}$ such that $\mathcal{A}=(\mathcal{A}_*)^*$, as a Banach space, for some Banach space $\mathcal{A}_*$, and such that $\mathcal{A}_*$ is a closed $\mathcal{A}-$submodule of $\mathcal{A}^*$; or equivalently, the multiplication on $\mathcal{A}$ is separately weak*-continuous. It should be remarked that the predual of a dual Banach algebra need not be unique, in general (see [6, 9]); so we usually point to the involved predual of a dual Banach algebra.

A functional $f\in\mathcal{A}^*$ is said to be *weakly almost periodic* if $\{f\cdot a : \|a\|\leq 1\}$ is relatively weakly compact in $\mathcal{A}^*$. We denote by $\mathcal{WAP}(\mathcal{A})$ the set of all weakly almost periodic elements of $\mathcal{A}^*$. It is easy to verify that, $\mathcal{WAP}(\mathcal{A})$ is a (norm) closed subspace of $\mathcal{A}^*$. For more about weakly almost periodic functionals, see [7].

It is known that the multiplication of a Banach algebra $\mathcal{A}$ has two natural but, in general, different extensions (called Arens products) to the second dual $\mathcal{A}^{**}$ each turning $\mathcal{A}^{**}$ into a Banach algebra. When these extensions are equal, $\mathcal{A}$ is said to be (Arens) regular. It can be verified that $\mathcal{A}$ is Arens regular if and only if $\mathcal{WAP}(\mathcal{A})=\mathcal{A}^*$. Further information for the Arens regularity of Banach algebras can be found in [6, 7].

$\mathcal{WAP}$-algebras, as a generalization of the Arens regular algebras, has been introduced and intensively studied in [8]. Indeed, a Banach algebra $\mathcal{A}$ for which

---

2010 *Mathematics Subject Classification.* 43A10, 43A20, 46H15, 46H25.

Key words and phrases. WAP-algebra, dual Banach algebra, Arens regularity, weak almost periodicity.

©2008 Universiteti i Prishtinës, Prishtinë, Kosovë.

Submitted xxxxx xxxxx. Published xxxxx xxxx.

Supported by the Center of Excellence for Mathematics at the Isfahan university.

the natural embedding of $\mathcal{A}$ into $\mathcal{WAP}(\mathcal{A})^*$ is bounded below, is called a $\mathcal{WAP}$-algebra. It has also known that $\mathcal{A}$ is a $\mathcal{WAP}$-algebra if and only if it admits an isomorphic representation on a reflexive Banach space. If $\mathcal{A}$ is a $\mathcal{WAP}$-algebra, then $\mathcal{WAP}(\mathcal{A})$ separate the points of $\mathcal{A}$ and so it is ω^*-dense in $\mathcal{A}^*$.

It can be readily verified that every dual Banach algebra, and every Arens regular Banach algebra, is a $\mathcal{WAP}$-algebra for comparison see [14]. Moreover group algebras are also always $\mathcal{WAP}$-algebras, however, they are neither dual Banach algebras, nor Arens regular in the case where the underlying group is not discrete, see [5, Corollary3.7] and [18, 14]. Ample information about $\mathcal{WAP}$-algebras with further details can be found in the impressive paper [8].

The paper is organized as follows. In section 2 we introduce some algebraic and topologic properties of $\mathcal{WAP}$-algebras. How to inherit the weakly almost periodicity of Banach algebras to the ideals, quotient spaces, direct sums and by homomorphisms are studied. We show that the second dual of a Banach algebra $\mathcal{A}$ is a $\mathcal{WAP}$-algebra, under each Arens products, if and only if $\mathcal{A}$ is Arens regular. In section 3 we deal with a locally compact foundation semigroup S, we show that the absolutely continuous semigroup measure algebra $M_a(S)$ is a $\mathcal{WAP}$-algebra if and only if the measure algebra $M_b(S)$ is so.

2. Some algebraic and topologic properties of $\mathcal{WAP}$-algebras

N. J. Young turns out a Banach algebra $\mathcal{A}$ admits an isometric representation on a reflexive Banach space if and only if the weakly almost periodic functionals of norm one determines the norm on $\mathcal{A}$. The following proposition provides a property qualification of $\mathcal{WAP}$-algebras.

Proposition 2.1. *Let $\mathcal{A}$ be a Banach algebra. Then the following statements are equivalent.*

(i) *$\mathcal{A}$ is a $\mathcal{WAP}$- algebra.*
(ii) *There is a reflexive Banach space E and a bounded below continuous representation $\pi : \mathcal{A} \to B(E)$.*
(iii) *There is a dual Banach algebra (or $\mathcal{WAP}$-algebra) $\mathfrak{B}$ and a bounded below continuous homomorphism $\phi : \mathcal{A} \to \mathfrak{B}$.*

Proof. (i)⇒(ii): The canonical map $\iota : \mathcal{A} \to \mathcal{WAP}(\mathcal{A})^*$ is bounded below and $\mathcal{WAP}(\mathcal{A})^*$ is dual Banach algebra. According to [8, Corollary3.8] there is a reflexive Banach space E such that $\pi : \mathcal{WAP}(\mathcal{A})^* \to B(E)$ is an isometric representation. So $\pi \circ \iota : \mathcal{A} \to B(E)$ is a bounded below continuous representation.

(ii)⇒ (iii) Trivial.

(iii)⇒ (i): Let $\iota : \mathcal{A} \to \mathcal{WAP}(\mathcal{A})^*$ be the canonical map. Suppose that $\mathfrak{B}$ is a dual Banach algebra and $\phi : \mathcal{A} \to \mathfrak{B}$ is a bounded below homomorphism, then $\phi^*(\mathcal{WAP}(\mathfrak{B})) \subseteq \mathcal{WAP}(\mathcal{A})$. Also:

$$\begin{aligned} \|\iota(a)\| &= \sup\{|f(a)| : f \in \mathcal{WAP}(\mathcal{A}) \quad \textit{and} \quad \|f\| \leq 1\} \\ &\geq \sup\{|\phi^*(g)(a)| : g \in \mathcal{WAP}(\mathfrak{B}) \quad \textit{and} \quad \|\phi\|\|g\| \leq 1\} \\ &= \frac{1}{\|\phi\|}\|\phi(a)\| \quad (\text{since } \mathfrak{B} \to \mathcal{WAP}(\mathfrak{B})^* \text{ is isometric}) \end{aligned}$$

So ι is bounded below. □

Corollary 2.2. *Let $\|.\|_1$ and $\|.\|_2$ be equivalent norm on algebra $\mathcal{A}$. Then $(\mathcal{A}, \|.\|_1)$ is a $\mathcal{WAP}$-algebra if and only if $(\mathcal{A}, \|.\|_2)$ is so.*

Proof. Since the identity map is bounded below, it is an immediate consequence of proposition 2.1 (iii). ☐

By above corollary, it is convenient to suppose a $\mathcal{WAP}$-algebra $\mathcal{A}$ has an isometric representation on reflexive space E. This can clearly be achieved by renorming $\mathcal{A}$ by $\|a\|_1 = \|\pi(a)\|, (a \in \mathcal{A})$ where π is a bounded below representation on E.

Corollary 2.3. *Let $\mathcal{A}$ be a $\mathcal{WAP}$-algebra and $\mathfrak{B}$ be a closed subalgebra of $\mathcal{A}$. Then $\mathfrak{B}$ is a $\mathcal{WAP}$-algebra.*

Proof. Since the inclusion map is bounded below, it is an immediate consequence of proposition 2.1 (iii). ☐

Proposition 2.4. *Let E be a Banach space. Then the following statements are equivalent:*

(i) *$B(E)$ is a dual Banach algebra.*
(ii) *$B(E)$ is a $\mathcal{WAP}$- algebra.*
(iii) *$E\hat{\otimes}E^* \subseteq \mathcal{WAP}(B(E))$.*
(iv) *E is a reflexive Banach space.*

Proof. (i)⇒ (ii) It is trivial.

(ii)⇒ (iv)If $E \neq \{0\}$ is not a reflexive Banach space, then $\mathcal{WAP}(F(E)) = \{0\}$ (see [18] Theorem 3). Thus $\mathcal{WAP}(B(E))$ can not separate the points of $F(E)$ and so $B(E)$. Thus it's not a $\mathcal{WAP}$-algebra.

(iv)⇒ (i)Let (T_α) be a net in $B(E)$ an $T, S \in B(E)$ such that $T_\alpha \xrightarrow{w^*} T$. Then for $\tau = \sum_{n=1}^{\infty} \xi_n \otimes \xi'_n$ in $E\hat{\otimes}E^*$, predual of $B(E)$, we have:

$$\begin{aligned} < ST_\alpha, \tau > &= \sum_{n=1}^{\infty} < ST_\alpha(\xi_n), \xi'_n > \\ &= \sum_{n=1}^{\infty} < T_\alpha(\xi_n), S^*(\xi'_n) > \longrightarrow \sum_{n=1}^{\infty} < T(\xi_n), S^*(\xi'_n) > \\ &= \sum_{n=1}^{\infty} < ST(\xi_n), \xi'_n > = < ST, \tau > \end{aligned}$$

And similarly $< T_\alpha S, \tau > \longrightarrow < TS, \tau >$. Hence the multiplication on $B(E)$ is separately w^*-continuous and $B(E)$ is a dual Banach algebra.

(iii)⇒ (iv)If $E \neq \{0\}$ is not a reflexive Banach space. Then $\mathcal{WAP}(B(E)) = \{0\}$([18, Theorem3]) and $E\hat{\otimes}E^* = \{0\}$. This is a contradiction.

(iv)⇒ (iii) See [18, Theorem 1]. ☐

The Proposition 2.1(iii)with Corollary 2.3 shows that a $\mathcal{WAP}$-algebra $\mathcal{A}$ is isomorphic to a closed subalgebra of a dual Banach algebra, so it is convenient to suppose that a $\mathcal{WAP}$-algebra is norm closed subalgebra of some dual Banach algebra. This can be achieved by redefining suitable norm.

A large class of $\mathcal{WAP}$-algebras are semisimple cummitative Bananach algebras. We refer to [13] for this kind of algebras.

Proposition 2.5. *Let $\mathcal{A}$ be a semisimple cummitative Bananach algebra, then $\mathcal{A}$ is $\mathcal{WAP}$-algebra.*

Proof. Since $\Delta(\mathcal{A}) \subseteq WAP(\mathcal{A})$ and $\Delta(\mathcal{A})$ seprerates the points of $\mathcal{A}$ and $||a||_1 = \sup\{|f(a)| : a \in \Delta(\mathcal{A})\}$ is a Banach norm on $\mathcal{A}$. We know that every Banach norm on semisimple Banach algebras are equivalent ([13]). Thus $\mathcal{A}$ is a WAP-algebra. □

Let G be a locally compact group and ω is a Borel-measurable weight function on it. Then the Fourier-Stieltjes algebra $B(G)$ and the weighted measure algebra $M_b(G,\omega)$ are dual Banach algebras, also the Fourier algebra $A(G)$ and $L^1(G,\omega)$ are closed ideals in them respectively. Hence all are $\mathcal{WAP}$-algebras.

Let $(\mathcal{A}, \mathcal{A}_*)$ be a dual Banach algebra and I is an ideal in $\mathcal{A}$, we define the annihilators:

$$I^{\perp} = \{f \in \mathcal{A}^* : f(i) = 0, \text{ for all } \quad i \in I\}, \quad {}^{\perp}I = \{f \in \mathcal{A}_* : f(i) = 0, \text{ for all } \quad i \in I\}.$$

It is easy to see that $({}^{\perp}I)^{\perp}$ is the ω^*-closure of I in $\mathcal{A}$.

Let $||a||_{\mathcal{WAP}(\mathcal{A})} = \sup\{|f(a)| : f \in \mathcal{WAP}(\mathcal{A}), ||f|| \leq 1\}$. Clearly $||.||_{\mathcal{WAP}(\mathcal{A})} \leq ||.||$. The following lemma show that whenever the seminorm $||.||_{\mathcal{WAP}(\mathcal{A})}$ could be a norm on $\mathcal{A}$.

Lemma 2.6. *Let $\mathcal{A}$ be a Banach algebra. Then the following conditions are equivalent.*

1. *$\mathcal{WAP}(\mathcal{A})$ separates the points of $\mathcal{A}$.*
2. *${}^{\perp}(\mathcal{WAP}(\mathcal{A})) = \{0\}$.*
3. *$\overline{\mathcal{WAP}(\mathcal{A})}^{\sigma(\mathcal{A}^*,\mathcal{A})} = \mathcal{A}^*$*
4. *$||.||_{\mathcal{WAP}(\mathcal{A})}$ is a norm on $\mathcal{A}$.*

Proof. Clearly (1) and (4)are equivalent.

(1)⇒(2). Let $0 \neq x \in \mathcal{A}$. Then there is a $f \in \mathcal{WAP}(\mathcal{A})$ such that $f(x) \neq 0$. Thus $x \notin {}^{\perp}(\mathcal{WAP}(\mathcal{A}))$, so ${}^{\perp}(\mathcal{WAP}(\mathcal{A})) = \{0\}$.

(2)⇒(3). $\mathcal{A}^* = 0^{\perp} = ({}^{\perp}(\mathcal{WAP}(\mathcal{A}))^{\perp} = \overline{\mathcal{WAP}(\mathcal{A})}^{\omega^*}$.

(3)⇒(1). Let $\overline{\mathcal{WAP}(\mathcal{A})}^{\omega^*} = \mathcal{A}^*$ then for $0 \neq x \in \mathcal{A}$ there is a $f \in \mathcal{A}^*$ such that $f(x) \neq 0$. Let $f_\alpha \in \mathcal{WAP}(\mathcal{A})$ convergent to f in ω^* topology then there is an index α such that $f_\alpha(x) \neq 0$ so $\mathcal{WAP}(\mathcal{A})$ separates the points of $\mathcal{A}$. □

By using [18, Theorem 1] and the previous lemma, the following is immediate.

Proposition 2.7. *The following statements are equivalent.*

1. *$\mathcal{A}$ is $\mathcal{WAP}$-algebra.*
2. *$\mathcal{WAP}(\mathcal{A})$ separates the points of $\mathcal{A}$ and $||.||_{\mathcal{WAP}(\mathcal{A})}$ and $||.||$ are equivalent.*

Theorem 2.8. *Let $\mathcal{A} = (\mathcal{A}_*)^*$ be a dual Banach algebra. Let I be a closed ideal in $\mathcal{A}$, then the following statements are equivalent.*

- (i) *$\frac{\mathcal{A}}{I}$ is a dual Banach algebra with respect to ${}^{\perp}I$.*
- (ii) *I is ω^*- closed ideal in $\mathcal{A}$.*

Proof. (ii)⇒(i) Let I be ω^*- closed ideal in $\mathcal{A}$. Then

$$\frac{\mathcal{A}}{I} = \frac{\mathcal{A}}{\overline{I}^{\omega^*}} = \frac{\mathcal{A}}{({}^{\perp}I)^{\perp}} = ({}^{\perp}I)^*$$

For $x \in \mathcal{A}$ and $f \in \mathcal{A}_*$ we define the module action of $\frac{\mathcal{A}}{I}$ over ${}^{\perp}I$ by $(x+I).f = x.f$ and $f.(x+I) = f.x$. It is routine to check that this module actions are well defined and $\frac{\mathcal{A}}{I}.({}^{\perp}I) \subseteq ({}^{\perp}I)$ and $({}^{\perp}I).\frac{\mathcal{A}}{I} \subseteq ({}^{\perp}I)$, so $\frac{\mathcal{A}}{I}$ is a dual Banach algebra with respect to ${}^{\perp}I$.

(ii)⇒(i). $\frac{\mathcal{A}}{I} = ({}^{\perp}I)^* = \frac{\mathcal{A}}{({}^{\perp}I)^{\perp}} = \frac{\mathcal{A}}{\overline{I}^{\omega^*}}$ thus $(\frac{\mathcal{A}}{I})^* = (\frac{\mathcal{A}}{\overline{I}^{\omega^*}})^*$. So ${}^{\perp}(I^{\perp}) ={}^{\perp} (\overline{I}^{\omega^*})^{\perp}$ and thus $I = \overline{I}^{\omega^*}$ □

In the next theorem $\mathcal{A}$ is a $\mathcal{WAP}$-algebra, so we suppose that $\mathcal{A}$ is a closed subalgebra of $\mathcal{WAP}(\mathcal{A})^*$.

Theorem 2.9. *Let $\mathcal{A}$ be a $\mathcal{WAP}$- algebra.*

- (i) *Let I be a $\sigma(\mathcal{A}, \mathcal{WAP}(\mathcal{A}))$ closed ideal in $\mathcal{A}$, then $\frac{\mathcal{A}}{I}$ is a $\mathcal{WAP}$- algebra.*
- (ii) *Let $\mathfrak{B}$ be a Banach algebra and $\varphi : \mathcal{A} \to \mathfrak{B}$ be an onto continuous homomorphism, then $\mathfrak{B}$ is a $\mathcal{WAP}$- algebra.*

Proof. (i)Let π be the canonical map from $\mathcal{A}$ onto $\mathcal{A}/I$. It is well known that $\pi^* : (\mathcal{A}/I)^* \longrightarrow \mathcal{A}^*$ is an isometric isomorphism onto it's rang $I^{\perp}$. We conclude from latter isometry that, $\mathcal{WAP}(\mathcal{A}/I) \cong \mathcal{WAP}(\mathcal{A}) \cap I^{\perp}$ is the predual of I in the dual Banach algebra $\mathcal{WAP}(\mathcal{A})^*$. I is a ω^*-closed ideal of $\mathcal{WAP}(\mathcal{A})^*$ with respect to both Arens products. So

$$\mathcal{WAP}(\mathcal{A})^*/I \cong \mathcal{WAP}(\mathcal{A}/I)^*$$

Since $\mathcal{A}$ is a $\mathcal{WAP}$-algebra, then $\mathcal{A}/I$ is a closed subalgebra of $\mathcal{WAP}(\mathcal{A})^*/I$ and so it is a $\mathcal{WAP}$-algebra.

(ii) Let $I = \ker(\varphi)$, then I is closed ideal in $\mathcal{A}$ and $\mathcal{A}/I \cong \mathfrak{B}$ isometrically isomorphic. Thus by (i) the Banach algebra $\mathfrak{B}$ is a $\mathcal{WAP}$-algebra. □

Let $\mathcal{A} \oplus \mathfrak{B}$ denote the direct sum of Banach algebras $\mathcal{A}$ and $\mathfrak{B}$, and endow it with coordinate wise operation.

Lemma 2.10.

- (i) *If $\mathcal{A}$ and $\mathfrak{B}$ are $\mathcal{WAP}$-algebras, then $\mathcal{A} \oplus \mathfrak{B}$ is $\mathcal{WAP}$-algebras as well.*
- (ii) *If $\mathcal{A}$ and $\mathfrak{B}$ are dual Banach algebras, then $\mathcal{A} \oplus \mathfrak{B}$ is dual Banach algebra as well.*
- (iii) *If $\mathcal{A}$ and $\mathfrak{B}$ are Arens regular Banach algebras, then $\mathcal{A} \oplus \mathfrak{B}$ is Arens regular as well.*

Corollary 2.11. *Let $\mathcal{A}^{\sharp}$ be the unitization of $\mathcal{A}$. Then:*

- (i) *$\mathcal{A}$ is a $\mathcal{WAP}$-algebras if and only if $\mathcal{A}^{\sharp}$ is $\mathcal{WAP}$-algebras as well.*
- (ii) *$\mathcal{A}$ is a dual Banach algebras if and only if $\mathcal{A}^{\sharp}$ is dual Banach algebra as well.*
- (iii) *$\mathcal{A}$ is Arens regular Banach algebras if and only if $\mathcal{A}^{\sharp}$ is Arens regular as well.*

Theorem 2.12. *Let $\mathcal{A}$ be a Banach algebra and I is a closed complemented ideal in $\mathcal{A}$. If I and $\frac{\mathcal{A}}{I}$ are dual Banach algebra [respectively, $\mathcal{WAP}$-algebra, Arens regular Banach algebra], then $\mathcal{A}$ is a dual Banach algebra [respectively, $\mathcal{WAP}$-algebra, Arens regular Banach algebra].*

Proof. (i): Let $I = E^*$ (respectively $\frac{\mathcal{A}}{I} = F^*$) where E (respectively F) is an I-bimodule (respectively $\frac{\mathcal{A}}{I}$-bimodule). Suppose that $\mathcal{A}_* := E \oplus F$. Then $\mathcal{A}_*^* \cong (E \oplus F)^* \cong E^* \oplus F^* \cong I \oplus \frac{\mathcal{A}}{I} \cong \mathcal{A}$. We can regard $\mathcal{A}_*$ as an $\mathcal{A}$-bimodule by

$$a.\mu := (x.\mu_1, y.\mu_2) \quad and \quad \mu.a := (\mu_1.x, \mu_2.y)$$

where $a = (x, y) \quad and \quad \mu = (\mu_1, \mu_2), x \in I, y \in \frac{\mathcal{A}}{I}, \mu_1 \in E, \mu_2 \in F$.

(ii):Suppose that I is a closed subalgebra of $\mathcal{WAP}(I)^*$ and $\frac{\mathcal{A}}{I}$ is closed subalgebra of $\mathcal{WAP}(\mathcal{A})^*/\bar{I}^{\omega^*} \cong \mathcal{WAP}(\mathcal{A}/I)^*$ then $\mathcal{A} \cong I \oplus \frac{\mathcal{A}}{I}$ is a closed subalgebra of $\mathcal{WAP}(I)^* \oplus \mathcal{WAP}(\mathcal{A})^*/\bar{I}^{\omega^*}$. So $\mathcal{A}$ is a $\mathcal{WAP}$-algebra.

(iii): Suppose that $\mathcal{WAP}(I) = I^*$ and $\mathcal{WAP}(\mathcal{A}/I) \cong (\mathcal{A}/I)^*$. Then $\mathcal{A}^* \cong (I \oplus \frac{\mathcal{A}}{I})^* \cong I^* \oplus (\frac{\mathcal{A}}{I})^* \cong \mathcal{WAP}(I) \oplus \mathcal{WAP}(\frac{\mathcal{A}}{I}) \cong \mathcal{WAP}(\mathcal{A})$. The last equality is a result of iterate limit argument. Let $F \in \mathcal{WAP}(I) \oplus \mathcal{WAP}(\frac{\mathcal{A}}{I})$ and (x_n, y_n) ,(a_m, b_m) are sequences in unit ball of $I \oplus \frac{\mathcal{A}}{I}$. Then $F = (f_1, f_2)$ such that $f_1 \in \mathcal{WAP}(I)$ and $f_2 \in \mathcal{WAP}(\frac{\mathcal{A}}{I})$. So that

$$\begin{aligned}
\lim_n \lim_m F(x_n, y_n).(a_m, b_m) &= (\lim_n \lim_m f_1(x_n a_m), \lim_n \lim_m f_2(y_n b_m)) \\
&= (\lim_m \lim_n f_1(x_n a_m), \lim_m \lim_n f_2(y_n b_m)) \\
&= \lim_m \lim_n F(x_n, y_n).(a_m, b_m)
\end{aligned}$$

and $F \in \mathcal{WAP}(\mathcal{A})$. ☐

Let X and Y be Banach spaces and $T : X^* \to Y^*$ be a bounded operator. Recall that the following conditions are equivalent:

(i) $T = S^*$, for some bounded operator $S : Y \to X$.
(ii) $T^*(Y) \subseteq X$.
(iii) T is ω^*-ω^*-continuous operator.

We now state the main result of this section.

Theorem 2.13. *Let $\mathcal{A}$ be a Banach algebra. Then the following statements are equivalent:*

(i) *$\mathcal{A}$ is a Arens regular Banach algebra.*
(ii) *$(\mathcal{A}^{**}, \Box)$ is a dual Banach algebra.*
(iii) *$(\mathcal{A}^{**}, \Diamond)$ is a dual Banach algebra.*
(iv) *$(\mathcal{A}^{**}, \Box)$ is a $\mathcal{WAP}$-algebra.*
(v) *$(\mathcal{A}^{**}, \Diamond)$ is a $\mathcal{WAP}$-algebra.*
(vi) *$\mathcal{A}$ is a $\mathcal{WAP}$- Banach algebra and $\mathcal{WAP}(\mathcal{A})$ is a ω^*-closed set in $\mathcal{A}^*$.*

Proof. If $\mathcal{A}$ is Arens regular, then by [1, Proposition 1.2] $\mathcal{A}^{**}$ is dual Banach algebra so(ii) and (iii) are equivalent. By [12, Theorem 4.10] there is a reflexive Banach space E such that $\mathcal{A}$ and also $\mathcal{A}^{**}$ are closed subalgebras of $B(E)$. Hence $\mathcal{A}^{**}$ is a $\mathcal{WAP}$-algebra. Thus (i)⇒ (ii)⇔ (iii)⇒ (iv) were established.

We show that if $\mathcal{A}^{**}$ with either Arens product has a representation on a reflexive Banach space then it is a representation with another Arens product. Thus the equality (iv)⇔ (v) was established. Let $\pi : (\mathcal{A}^{**}, \Box) \to B(E)$ be an isometric representation on a reflexive Banach space E (See corollary2.2 and explanations after it). By [8, Proposition 4.2] π is ω^*-ω^*-continuous. Let $\Phi, \Psi \in \mathcal{A}^{**}$ and (a_α) and (b_β) be nets in $\mathcal{A}$ that ω^*-convergening to Φ and Ψ, respectively. Then for each $\tau \in E \hat{\otimes} E^*$

$$\begin{aligned}
< \tau, \pi(\Phi \Box \Psi) > &= \lim_\alpha \lim_\beta < \tau, \pi(a_\alpha) \circ \pi(b_\beta) > \\
&= \lim_\beta \lim_\alpha < \tau, \pi(a_\alpha) \circ \pi(b_\beta) > \qquad (\text{since } \tau \in \mathcal{WAP}(B(E))) \\
&= < \tau, \pi(\Phi \Diamond \Psi) >
\end{aligned}$$

So $\pi : (\mathcal{A}^{**}, \Diamond) \to B(E)$ is an isometric representation. Similarly for another Arens product.

(v)$\Rightarrow$ (i)Let $(\mathcal{A}^{**}, \Diamond)$ is a $\mathcal{WAP}$-algebra. Then there is an isometric representation $\pi : (\mathcal{A}^{**}, \Diamond) \to B(E)$, for some reflexive Banach space E. Hence by above equalities,

$$\pi(\Psi \Diamond \Phi) = \pi(\Psi) \circ \pi(\Phi) = \pi(\Psi \Box \Phi) \quad (\Phi, \Psi \in \mathcal{A}^{**})$$

Since π is injective, we conclude $\mathcal{A}$ is Arerns regular Banach algebra.

(vi)$\Rightarrow$ (i) Let $\mathcal{A}$ be a $\mathcal{WAP}$-algebra and $\mathcal{WAP}(\mathcal{A})$ be a ω^*-closed set in $\mathcal{A}^*$. By lemma 2.6 $\overline{\mathcal{WAP}(\mathcal{A})}^{\sigma(\mathcal{A}^*,\mathcal{A})} = \mathcal{A}^* = \mathcal{WAP}(\mathcal{A})$.

(i)$\Rightarrow$ (vi) Let $\mathcal{A}$ be a a regular Banach algebra. Since $\mathcal{A}$ is a closed subset of $\mathcal{A}^{**}$, $\mathcal{A}$ is a $\mathcal{WAP}$-algebra. and $\overline{\mathcal{WAP}(\mathcal{A})}^{\sigma(\mathcal{A}^*,\mathcal{A})} = \mathcal{A}^* = \mathcal{WAP}(\mathcal{A})$. So $\mathcal{WAP}(\mathcal{A})$ is a ω^*-closed subset of $\mathcal{A}^*$. □

Proposition 2.14. *$\mathcal{A}^{**}$ is a Arens regular Banach algebra if and only if $\mathcal{A}$ is a Arens regular Banach algebra and $\mathcal{WAP}(\mathcal{A}^{**}) = \mathcal{WAP}(\mathcal{A})^{**}$.*

Proof. Let $\mathcal{A}^{**}$ be a Arens regular Banach algebra, then $\mathcal{WAP}(\mathcal{A}^{**}) = \mathcal{A}^{***}$. Also $\mathcal{A}$ is a closed subalgebra of $\mathcal{A}^{**}$ so it is Arens regular Banach algebra. Thus $\mathcal{WAP}(\mathcal{A}) = \mathcal{A}^*$ and so $\mathcal{WAP}(\mathcal{A}^{**}) = \mathcal{WAP}(\mathcal{A})^{**}$.

Conversely, let $\mathcal{A}$ be a Arens regular Banach algebra then $\mathcal{WAP}(\mathcal{A}) = \mathcal{A}^*$. Thus $\mathcal{WAP}(\mathcal{A}^{**}) = \mathcal{WAP}(\mathcal{A})^{**} = \mathcal{A}^{***}$, and so $\mathcal{A}^{**}$ is Arens regular. □

There exists a Banach algebra $\mathcal{A}$ such that $\mathcal{A}$ is Arens regular, but $\mathcal{A}^{**}$ is not Arens regular (see [7]).

Examples 2.15.
1. *Let $S = (\mathbb{N}, \min)$. Then $\ell_1(S)$ is a $\mathcal{WAP}$-algebra, but the second dual isn't so.*
2. *Let $S = (\mathbb{N}, \max)$. Then $\ell_1(S)$ is a dual Banach algebra, but the second dual isn't so.*
3. *Let S be infinite zero semigroup, then $\ell_1(S)^{**}$ is a dual Banch algebra but $\ell_1(S)$ isn't so.*

3. ABSOLUTELY SEMIGROUP MEASURE ALGEBRAS

Following [4], a semitopological semigroup is a semigroup S equipped with a Hausdorff topology under which the multiplication of S is separately continuous. If the multiplication of S is jointly continuous then S is said to be a topological semigroup. We write $\ell^\infty(S)$ for the commutative algebras of all bounded complex-valued functions on S. In the case where S is locally compact we also write $C(S)$ and $C_0(S)$ for the closed subalgebras of $\ell_\infty(S)$ consist of continuous elements and continuous elements which vanish at infinity, respectively. We also denote the space of all *weakly almost periodic* functions on S by $\mathcal{WAP}(S)$ which is defined by

$$\mathcal{WAP}(S) = \{f \in C(S) : \{R_s f : s \in S\} \text{ is relatively weakly compact in } \quad C(S)\},$$

where $R_s f(t) = f(ts)$, $(s, t \in S)$. Then $\mathcal{WAP}(S)$ is a closed subalgebra of $C(S)$. Many other properties of $\mathcal{WAP}(S)$ and its inclusion relations among other function algebras are completely explored in [4].

Let $M_b(S)$ denotes the Banach space of all bounded complex regular Borel measures on S with the total variation norm. Then $M_b(S)$ with the convolution multiplication $*$ defined by the equation

$$\langle \mu * \nu, f\rangle = \int_S \int_S f(xy) d\mu(x) d\nu(y) \quad (f \in C_0(S))$$

is a convolution measure algebra as the dual of $C_0(S)$.

Let $\mu \in M_b(S)$ and let $L^\infty(|\mu|)$ be the Banach space of all bounded Borel-measurable functions on S with essential supremum norm,

$$||f||_\mu = \inf\{\alpha \geq 0 : \{x \in S : |f(x)| > \alpha\} \quad \text{is} \quad |\mu| - \text{null}\}.$$

Consider the product linear space $\Pi\{L^\infty(|\mu|) : \mu \in M_b(S)\}$. An element $f = (f_\mu)$ in this product is called a generalized function on S if the following conditions are satisfied:

(1) $||f||_\infty = \sup\{||f||_\mu : \mu \in M_b(S)\}$ is finite.
(2) if $\mu, \nu \in M_b(S)$ and $|\mu| \ll |\nu|$, then $f_\mu = f_\nu$, $|\mu|$-a.e.

Let $GL(S)$ denote the linear subspace of all generalized functions on S. Then $M_b(S)^*$ is isometrically isomorphic to $GL(S)$, see [15]. We identify $C_0(S)$ with its image by canonical map $\iota : C_0(S) \to GL(S)$. The w^*-topology on $GL(S)$ is $\sigma(GL(S), M_b(S))$. Let $\mathcal{A}$ be a convolution measure algebra in $M_b(S)$, i.e. a norm-closed solid subalgebra of $M_b(S)$. In other words, for all $\mu \in M_b(S)$ and $\nu \in \mathcal{A}$ such that $|\mu| \ll |\nu|$, we have $\mu \in \mathcal{A}$. The space of all measures $\mu \in M_b(S)$ for which the maps $x \mapsto \delta_x * |\mu|$ and $x \mapsto |\mu| * \delta_x$ from S into $M_b(S)$ are weakly continuous is denoted by $M_a(S)$, where δ_x denotes the Dirac measure at x. Then $M_a(S)$ is a closed two-sided L-ideal of $M_b(S)$; see Baker and Baker [3] or Dzinotyiweyi [11]. The locally compact semigroup S is called foundation if , $F_a(S)$, the closure of the set

$$\cup\{supp(\mu) : \mu \in M_a(S)\}$$

coincide with S .

Let F and K be nonempty subsets of a semigroup S and $s \in S$. We put

$$s^{-1}F = \{t \in S : st \in F\}, \text{ and } \quad Fs^{-1} = \{t \in S : ts \in F\}$$

and we also write $s^{-1}t$ for the set $s^{-1}\{t\}$, FK^{-1} for $\cup\{Fs^{-1} : s \in K\}$ and $K^{-1}F$ for $\cup\{s^{-1}F : s \in K\}$.

The authors and H.R. Ebrahimi- Vishki in [14], for a locally compact topological semigroup S showed that $M_b(S)$ is a $\mathcal{WAP}$-algebra if and only if the canonical mapping $R : S \longrightarrow S^{wap}$ is one to one, where S^{wap} is the weakly almost periodic compactification of S. In this section we show that for a foundation semigroup S, this is equivalent to $M_a(S)$ is a $\mathcal{WAP}$-algebra.

The following theorem is the main result of this section.

Theorem 3.1. *Let S be a foundation locally compact topological semigroup with identity. Then $M_a(S)$ is a $\mathcal{WAP}$-algebra if and only if $M_b(S)$ is so.*

Proof. If $M_b(S)$ is a $\mathcal{WAP}$-algebra then $M_a(S)$ is a closed subalgebra in it, so $M_a(S)$ is a $\mathcal{WAP}$-algebra.

As the proof of lemma 1.8 [11, p.16] for all $x_1, x_2 \in S$ with $x_1 \neq x_2$ there is $\mu \in M_a(S)$ such that

$$\delta_{x_1} * \mu \neq \delta_{x_2} * \mu.$$

Let $M_a(S)$ is a $\mathcal{WAP}$-algebra, then for all $\mu \in M_a(S)$ we have $||\mu|| = \sup\{|\langle f, \mu\rangle| : ||f|| \leq 1, f \in \mathcal{WAP}(M_a(S))\}$ and by [15, Theorem3.7] $\mathcal{WAP}(S) = \mathcal{WAP}(M_a(S))$, so $||\mu|| = \sup\{|\langle f, \mu\rangle| : ||f|| \leq 1, f \in \mathcal{WAP}(S)\}$. Thus there is a $f \in \mathcal{WAP}(S)$ such that

$$\langle \mu, R_{x_1} f\rangle = \langle \delta_{x_1} * \mu, f\rangle \neq \langle \delta_{x_2} * \mu, f\rangle = \langle \mu, R_{x_2} f\rangle$$

Thus $R_{x_1} \neq R_{x_2}$. This means the canonical mapping $R : S \longrightarrow S^{wap}$ is one to one, where S^{wap} is the weakly almost periodic compactification of S. By [14, Theorem 3.1] $M_b(S)$ is a $\mathcal{WAP}$-algebra. □

By using [14, Theorem3.1] and the previous Theorem, the following is immediate.

Corollary 3.2. *Let S be a foundation semigroup. The following statements are equivalent.*

1. *$M_a(S)$ is $\mathcal{WAP}$-algebra.*
2. *$\ell_1(S)$ is a $\mathcal{WAP}$-algebra.*
3. *$M_b(S)$ is a $\mathcal{WAP}$-algebra.*

By using [16, Corollary8] and Theorem2.13, the following is immediate.

Corollary 3.3. *Let S be a foundation semigroup. The following statements are equivalent.*

1. *$M_a(S)^{**}$ is $\mathcal{WAP}$-algebra.*
2. *$\ell_1(S)^{**}$ is a $\mathcal{WAP}$-algebra.*
3. *$M_b(S)^{**}$ is a $\mathcal{WAP}$-algebra.*

The next example shows that in Theorem3.1 the condition S be foundation semigroup can not be omitted.

Examples 3.4. (a) *Let S be a locally compact semigroup. If $C_0(S) \subseteq \mathcal{WAP}(S)$ then $M_a(S)$ is a $\mathcal{WAP}$-algebra.*

(b) *Let S be a compact semigroup, then $M_a(S)$ is a $\mathcal{WAP}$-algebra.*

(c) *Let $S = \mathbb{R}^2$ and $(x, y).(\alpha, \beta) = (\alpha x, \alpha y + \beta)$ with the standard Euclidean topology. Then $M_a(S) = \{0\}$ and so S is not a foundation semigroup. In fact, let $x_n = (0, 1 + \frac{1}{n})$, $x_0 = (0, 1)$, $K = [-1, 1] \times [-1, 1]$ and $\mu \in M_a^r(S)$. Then $Kx_n^{-1} = \emptyset$ and $Kx_0^{-1} = S$. Then by continuity of μ, we have $0 = |\mu|(Kx_n^{-1}) \to |\mu|(Kx_0^{-1}) = ||\mu||$. So $M_a(S) = \{0\}$.*

Let $f \in C_0(S)$ and $f(0, c) \neq 0$ for some $c \in \mathbb{R}$. Suppose $a_n = (0, n)$ and $b_m = (1, \gamma_m)$ where γ_m is on the line $y = -nx + c$. Then

$$\lim_n \lim_m f(a_n b_m) = f(0, c) \neq \lim_m \lim_n f(a_n b_m) = 0$$

So by examining the iterate limits we certify that $f \notin \mathcal{WAP}(S)$. Thus $\mathcal{WAP}(S)$ does not separate the points of Y and so $M_b(S)$. This show that the map $R : S \to S^{wap}$ is not one to one. Therefore $M_b(S)$ isn't a $\mathcal{WAP}$-algebra (see [14] also) but clearly $M_a(S)$ is a $\mathcal{WAP}$- algebra.

Acknowledgments. This research was supported by the Center of Excellence for Mathematics at the Isfahan university.

References

[1] M. Abolghasemi, A. Rejali, and H.R. Ebrahimi Vishki, *Weighted semigroup algebras as dual Banach algebras,* arXiv:0808.1404 [math.FA].

[2] F. Abtahi, B. Khodsiani, and A. Rejali, *Arens regularity of inverse semigroup algebras* , Bull. Iranian Math. Soc. Vol. 40 (2014), No. 6, pp. 1527-1538.

[3] A.C. Baker, J.W. Baker, *Algebras of measures on a locally compact semigroups,* **1**, J.London Math., no.1 (1969), 249-259.

[4] J.F. Berglund, H.D. Junghenn and P. Milnes, *Analysis on Semigroups*, Wiley, New York, (1989).

[5] R.B. Burckel, *Weakly Almost Periodic Functions on Semigroups,* Gordon and Breach, New york,(1970).

[6] H.G. Dales, *Banach Algebras and Automatic Continuity,* Clarendon Press, Oxford, 2000.

[7] H.G. Dales, A.T.M. Lau, *The Second Dual of Beurling Algebras,* Mem. Amer. Math. Soc. **177** (2005), no. 836.

[8] M. Daws, *Dual Banach algebras: Representation and injectivity,* Studia Math, **178**(2007), 231-275.

[9] M. Daws, H.L. Pham and S. White, *Conditions implying the uniqeness of the weak$^*-$topology on certain group algebras*, Houston J. Math. **35** (2009), no. 1, 253-276.

[10] H.A.M. Dzinotyiweyi, *Weakly almost periodic functions and the irregularity of multiplication in semiogroup algebras,* Math. Scand.**46**, (1980), 157-172.

[11] H.A.M. Dzinotyiweyi, *The Analogue of the Group Algebra for Topological Semigroups,* Reserch notes in mathematics, Pitman Publishing,(1984).

[12] S. Kaijser, *On Banach modules I*, Math. Proc. Cambridge Philos. Soc. **90** (1981).

[13] E.Kaniuth, *A Course in Commutative Banach Algebras* Springer New York 2008.

[14] H.R. Ebrahimi Vishki, B. Khodsiani and A. Rejali, *Weighted semigroup measure algebra as a $\mathcal{WAP}$-algebra,* U.P.B. Scientific Bulletin Accepted for publication.

[15] M. Lashkarizadeh Bami, *Function algebras on weighted topological semigroups,* Math. Japonica **47**, no.2 (1998), 217-227.

[16] A. Rejali, *The Arens regularity of weighted semigroup algebras,* Sci. Math. Japon, **60**, no.1 (2004), 129-137.

[17] A. Ulger, *Continuity of weakly almost periodic functionals on* $L^1(G)$, Quart. J. Math. Oxford(2), 37 (1986), 495-497.

[18] N.J.Young, Periodicity of functionals and representations of normed algebras on reflexive spaces., Proc. Edinburgh Math. Soc.(2)**20** (1976/77), 99-120.

Bahram Khodsiani
Department of Mathematics, Isfahan University , IRAN
E-mail address: `b_khodsiani@sci.ui.ac.ir`

Ali Rejali
Department of Mathematics, Isfahan University , IRAN.

E-mail address: `rejali@sci.ui.ac.ir`